

Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 1)

Vaccine Requirement	Child Care Facility, Preschool, Early Childhood Pre-Kindergarten Programs	Kindergarten through 12 th Grade		Minimum Intervals Allowed Between Doses and <i>Other Options</i> for Proof of Immunity
		First Entry into School (Kindergarten or First Grade)	Other Grades	
DTP/DTaP/ or Tdap, Td (Diphtheria, Tetanus, Pertussis)	Three doses by 1 year of age One additional booster dose by 2 nd birthday	Four or more doses of DTP/DTaP with the last dose qualifying as a booster and received on or after the 4 th birthday	Three or more doses of DTP/DTaP or Td; with the last dose qualifying as a booster if received on or after the 4 th birthday For Students entering 6 th thru 12 th grades: 1 dose of Tdap	Minimum interval between series doses: 4 weeks (28 days) Between series and booster: 6 months <i>No proof of immunity allowed</i>
Polio	Two doses by 1 year of age. One additional dose by 2 nd birthday	Four or more doses of the same type of Polio vaccine with the last dose qualifying as a booster and received on or after the 4 th birthday. (progressive requirement)	Three or more doses of Polio with the last dose qualifying as a booster and received on or after the 4 th birthday. *If the series is given in any combination of polio vaccine types, 4 or more doses are required with the last being a booster on or after the 4 th birthday.	Minimum interval between series doses: 4 weeks (28 days) For Grade K: 6 month interval between 3 dose series and booster; booster must be on or after 4 th birthday <i>No proof of immunity allowed</i>
Measles	One dose on or after the 1 st birthday but prior to 24 months of age	Two doses of Measles Vaccine, the 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.		<i>Laboratory evidence of measles immunity OR Certified physician verification* of measles disease by date of illness</i> *Cases diagnosed after 7/1/2002 must include lab evidence of infection
Rubella	One dose on or after the 1 st birthday but prior to 24 months of age	Two doses of Rubella Vaccine, the 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.		<i>Laboratory evidence of rubella immunity</i> History of disease is not acceptable proof of immunity to rubella
Mumps	One dose on or after the 1 st birthday but prior to 24 months of age	Two doses of Mumps Vaccine, the 1 st dose must have been received on after the 1 st birthday and the second dose no less than 4 weeks (28 days) later.		<i>Laboratory evidence of mumps immunity OR Certified physician verification of mumps disease by date of illness</i>
<i>Haemophilus influenzae</i> type b	Refer to Hib ACIP recs for <u>schedule for series</u> Children 24-59 mos. without series must have one dose after 15 mos. of age	Not required after the 5 th birthday (60 months of age)		Refer to ACIP Hib vaccination schedule <i>No proof of immunity allowed</i>
Invasive Pneumococcal Disease	Refer to PCV ACIP recs for <u>schedule for series</u> Children 24-59 mos. without series must have one dose after 24 months of age	Not required after the 5 th birthday (60 months of age)		Refer to ACIP PCV vaccination schedule <i>No proof of immunity allowed</i>

Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2016 (Page 2)

Vaccine Requirement	Child Care Facility, Preschool, Early Childhood Pre-Kindergarten Programs	Kindergarten through 12 th Grade		Minimum Intervals Allowed Between Doses and Other Options for Proof of Immunity
		First Entry into School (Kindergarten or First Grade)	All Other Grades	
Hepatitis B	<p>Three doses for all children</p> <p>Third dose must have been administered on or after 6 months of age (168 days)</p>	No Requirements	<p>For Students entering grades 6 thru 12:</p> <p>Three doses hepatitis B vaccine administered at recommended intervals</p>	<p>Minimum intervals between doses: 1 & 2- at least 4 weeks (28 days) 2 & 3 - at least 2 months (56 days) 1 & 3 - at least 4 months (112 days) <i>Laboratory evidence of prior or current infection</i></p>
Varicella (progressive requirement)	One dose on or after 1 st birthday	<p>Two doses of Varicella Vaccine, for Students entering Kindergarten or first grade</p> <p>The 1st dose must have been received on after the 1st birthday and the second dose no less than 4 weeks (28 days) later.</p>	<p>One dose of Varicella on or after the 1st birthday for Students entering grades 3, 4, 5, & 12</p> <p>Two doses of Varicella Vaccine for Students entering grades 2, 6, 7, 8, 9, 10, & 11</p>	<p>Minimum intervals for administration: The 1st dose must have been received on after the 1st birthday and the second dose no less than 4 weeks (28 days) later. <i>Statement from physician or health care provider verifying disease history OR Laboratory evidence of varicella immunity</i></p>
Meningococcal Conjugate Vaccine (progressive requirement)	No Requirements	No Requirements	<p>Applies to Students entering grades 6, 7, or 12 beginning 2016-2017 school year</p> <p>1 dose of Meningococcal vaccine for entry to grade 6 or 7</p> <p>2 doses of Meningococcal vaccine at entry to 12th grade</p>	<p>Minimum intervals for administration: The first dose received on or after the 10th birthday; second dose at least eight weeks after the 1st dose. Only one dose is required if the first dose was received at 16 years of age or older. <i>No proof of immunity allowed.</i></p>

Source: Child Health Examination Code/Part 665

Prepared by: Illinois Department of Public Health Immunization Section, January 2016