

KEWANEE COMMUNITY UNIT SCHOOL DISTRICT 229 HEALTH REQUIREMENTS

PRE-K:

Physical Exam

Lead Level (must have Blood Lead Level, not just a screen/assessment)

(Recommended: 2 Lead Levels – one year apart)

DTP/DTaP/Td (4 or more doses, at intervals of no less than 4 weeks apart)

Polio (3 or more doses, at intervals of no less than 4 weeks apart)

Measles (1 dose, administered on or after 1st birthday – or had disease)

Mumps (1 dose, administered on or after 1st birthday – or had disease)

Rubella (1 dose, administered on or after 1st birthday – or had disease)

Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)

Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months; 3rd/last dose – child must be 6 months of age or older)

Hib (children 24-59 months of age who have not received a primary series according to the Hib schedule, must show proof of receiving 1 dose at 15 months of age or older)

Pneumococcal Vaccine (children 24-59 months of age who have not received a primary series according to the recommended schedule, must show proof of receiving 1 dose of pneumococcal vaccine. Any child who has reached his or her 5th birthday shall not be required to provide proof of immunization with the vaccine).

KINDERGARTEN:

Physical Exam

Eye Exam

Dental Exam

Lead Level (must have Blood Lead Level, not just a screen/assessment)

(Recommended: 2 Lead Levels – one year apart)

DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)

Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)

Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Varicella (2 doses, 1st dose administered on or after 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease – need month & year of disease)

FIRST GRADE:

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)
- Any other physical and/or immunizations that are not complete

SECOND GRADE:

- Dental Exam
- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)
- Any other physical and/or immunizations that are not complete

THIRD GRADE (cont'd on next page):

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

THIRD GRADE (cont'd):

Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)

Any other physical and/or immunizations that are not complete

FOURTH GRADE:

DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)

Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)

Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)

Any other physical and/or immunizations that are not complete

FIFTH GRADE:

DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)

Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)

Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)

Any other physical and/or immunizations that are not complete

SIXTH GRADE (cont'd on next page):

Physical Exam

Dental Exam

DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)

TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose.

SIXTH GRADE (cont'd):

- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (2 doses, 1st dose administered on or after 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease – need month & year of disease)
- Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)
- Meningococcal Conjugate Vaccine (MCV – 1 dose on or after 11th birthday)
- Any other immunizations that are not complete

SEVENTH GRADE:

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose.
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)
- Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)
- Any other physical and/or immunizations that are not complete

EIGHTH GRADE (cont'd on next page):

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose

EIGHTH GRADE (cont'd):

- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)
- Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)
- Any other physical and/or immunizations that are not complete

NINTH GRADE:

- Physical Exam
- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose.
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (2 doses, 1st dose administered on or after 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease – need month & year of disease)
- Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)
- Any other immunizations that are not complete

TENTH GRADE (cont'd on next page):

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose.

TENTH GRADE (cont'd):

- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)
- Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)
- Any other physical and/or immunizations that are not complete

ELEVENTH GRADE:

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose.
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)
- Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)
- Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)
- Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)
- Any other physical and/or immunizations that are not complete

TWELFTH GRADE (cont'd on next page):

- DTP/DTaP/Td (4 or more doses, with last dose administered on or after the 4th birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose)
- TDaP Vaccine – must have one dose of TDaP vaccine regardless of the interval since the last DTaP, DT or Td dose.
- Polio (3 or more doses, at intervals of no less than 4 weeks apart, and the last dose having been received on or after the 4th birthday)

TWELFTH GRADE (cont'd):

Measles (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Mumps (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Rubella (2 doses, 1st dose administered on or after the 1st birthday, 2nd dose no less than 4 weeks after 1st dose – or had disease)

Varicella (1 dose, administered on or after 1st birthday or confirm past disease history – need month & year of disease)

Hepatitis B (3 doses, interval of at least 4 weeks between 1st & 2nd doses; interval between 2nd & 3rd doses must be at least 2 months; interval between 1st and 3rd doses must be at least 4 months)

Meningococcal Conjugate Vaccine (MCV – 2 doses – 1st dose received on or after the 11th birthday, and the 2nd dose received on or after 16th birthday, interval must be at least 8 weeks after the 1st dose. If the 1st dose is received at 16 years of age or older, only 1 dose is required)

Any other physical and/or immunizations that are not complete

STUDENTS ENTERING THE SCHOOL DISTRICT FOR THE FIRST TIME OR FROM OUT OF STATE MUST HAVE AN ILLINOIS PHYSICAL EXAM, DENTAL EXAM AND EYE EXAM COMPLETED (Waivers are available for the Dental Exam and Eye Exam if needed)

ANY OTHER PHYSICAL AND/OR IMMUNIZATIONS THAT ARE NOT COMPLETE NEED TO BE COMPLETED AS IDENTIFIED.

Developed: 5/05

Revised: 7/07, 4/08, 7/08, 7/09, 7/10, 7/11, 6/12, 7/13, 6/14, 5/15

Mary McIntire, RN, BSN