

Table 1: Considerations for Persons with Systemic Signs and Symptoms Pre- and Post- COVID-19 Vaccination

Recommended Infection Prevention and Control Strategies	Status of Individual with Systemic Signs and Symptoms of COVID-19 Infection <sup>1, 2</sup>				
	Unvaccinated	Received COVID-19 Vaccination in the prior 3 days (including day of vaccination, which is considered Day 1) <sup>7</sup>			Partially <sup>4</sup> or Fully <sup>5</sup> Vaccinated
		Symptoms unlikely to be from COVID-19 vaccination <sup>1</sup> (e.g., cough, shortness of breath, rhinorrhea, loss of taste/smell)	Symptoms <i>may</i> be vaccine related, possible COVID-19, or other unrelated illness <sup>2</sup> (e.g., chills, headache, myalgia, arthralgia, fatigue)		
<b>Exclude from School</b>	YES	YES	NO- if afebrile <sup>3</sup> and symptoms <sup>2</sup> occur within the first 48 hours and feels well enough to work	YES – if fever of 100.4F or greater or if symptoms persist for more than 2 days.	YES
<b>Refer for Clinical Evaluation</b>	YES Refer to <b>Row D</b> in <u>Exclusion Guidance Decision Tree</u> .	YES	YES – if symptoms do not improve and persist for more than 2 days. Not Recommended - if symptoms resolve within 2 days.		YES Refer to <b>Row D</b> in <u>Exclusion Guidance Decision Tree</u> .
<b>Perform COVID-19 Test</b>  (Healthcare Provider may decide to test based on clinical assessment.)	YES  NOT Recommended – if less than 90 days (3 months) from last positive test.	YES  Not Recommended – if less than 90 days (3 months) from last positive test	YES – if symptoms do not improve and persist for more than 2 days.  Not Recommended if symptoms resolve within 2 days  Not Recommended – if less than 90 days (3 months) from last positive test		NOT Recommended – if less than 90 days (3 months) from last positive test.
If SARS-CoV-2 test <b>Positive</b> : Refer to <b>Row A</b> in <u>Exclusion Guidance Decision Tree</u> . If SARS-CoV-2 test <b>Negative</b> <sup>6</sup> : Refer to <b>Row B</b> in <u>Exclusion Guidance Decision Tree</u> .					
<b>Place in Isolation</b>	YES – if SARS-CoV-2 test <b>Positive</b> : Stay home at least ten calendar days from onset of symptoms AND for 24 hours with no fever (without fever-reducing medication) AND improvement of symptoms.		NO- if afebrile <sup>3</sup> and symptoms occur within the first 48 hours and feels well enough to work.  Recommended - if symptoms persist for more than 2 days and pending test results.  YES - if SARS-CoV-2 test <b>Positive</b> .		YES – if SARS-CoV-2 test <b>Positive</b>

**Table 2: Considerations for Asymptomatic Persons Pre- and Post- COVID-19 Vaccination**

Recommended Infection Prevention and Control Strategies	Status of Asymptomatic Individual		
	Unvaccinated	Received COVID-19 Vaccination in the prior 3 days (including day of vaccination, which is considered Day 1) or is Partially <sup>4</sup> Vaccinated	Fully <sup>5</sup> Vaccinated
Place in Quarantine if named as a Close Contact to a Known Case of COVID-19	<p>YES – if no history of previous positive test for SARS-CoV-2; or if greater than 90 days (3 months) from last positive test.</p> <p>Not Recommended – if less than 90 days (3 months) from last positive test.</p>	<p>YES – if no history of previous positive test for SARS-CoV-2; or if greater than 90 days (3 months) from last positive test.</p> <p>Not Recommended – if less than 90 days (3 months) from last positive test.</p>	<p>NO -if they meet <u>all</u> the following criteria:</p> <ul style="list-style-type: none"> <li>• Are fully vaccinated (i.e., ≥2 weeks following receipt of the second dose in a 2-dose series, or ≥2 weeks following receipt of one dose of a single-dose vaccine)</li> <li>• Have remained asymptomatic since the current COVID-19 exposure</li> </ul> <p>YES – if they do not meet all three of the above criteria.</p>

1 Presence of ANY systemic signs and symptoms consistent with COVID-19 infection (e.g., cough, shortness of breath, rhinorrhea, sore throat, loss of taste or smell) or another infectious etiology (e.g., influenza) that are not typical for post-vaccination signs and symptoms.

2 Signs and symptoms *that may be* from either COVID-19 vaccination, SARS-CoV-2 infection, or another infectious etiology (e.g., fever of 100.4 or higher, fatigue, headache, chills, myalgia, arthralgia). **For symptomatic persons who are close contacts to a COVID-19 case, isolate, send home, and refer for testing and clinical evaluation.**

3 Must be afebrile for at least 24 hours to return to work.

4 Person receiving the first dose of a two-dose vaccine series OR person receiving one dose of a single-dose vaccine OR the second dose in a 2-dose series AND is <2 weeks following receipt of the last dose in the series.

5 Person receiving one dose of a single-dose vaccine OR the second dose in a 2-dose series AND is ≥2 weeks following receipt of the last dose in the series.

6 If performed, a negative SARS-CoV-2 antigen test in a person who has signs and symptoms that are typical for COVID-19 infection should be confirmed by SARS-CoV-2 nucleic acid amplification test (NAAT), e.g., RT-PCR test.

7 Individuals who are close contacts to COVID-19 case should follow quarantine guidelines; symptomatic close contacts should be tested for COVID-19.

Resources:

[Interim Public Health Recommendations for Fully Vaccinated People](#)

[Public health recommendations for vaccinated persons](#)

[Quarantine for those with natural immunity](#)