

Customer Notice of Lead and Copper Results in Drinking Water (NTNC/OTM)

As a public water system we are required to test our tap water for lead and copper corrosion products from actual drinking water taps and notify participants of their test results. Listed below are test results of samples collected this year

Public Water Supply Name	Fairview School		
PWSID.	44305877	County:	Oconto

Thank you for participating in our drinking water lead and copper monitoring program. The results of the lead and copper sample collected at your location are in the table below.

Address / Location of Sample	Sample Date	Lead (ug/L)	Copper (ug/L)
2840 HWY 32 - Krakow			
Water Cooler South End	09/10/19	<1.4	<20
Room 2	09/10/19	<1.4	<20
Kitchen Tap	09/10/19	<1.4	<20
Room 22	09/10/19	<1.4	<20
Water Cooler North End	09/10/19	<1.4	<20

Contaminant	Action Level (parts per billion or ppb)	Maximum Contaminant Level Goal (ppb)
Lead	15	0
Copper	1300	1300

Definition of terms

ug/L: micrograms per liter. This is equivalent to one part-per-billion or *ppb* .

Action Level: The concentration of lead or copper which, if exceeded in greater than ten percent of the total number of samples collected by a public water system, triggers one or more of the actions specified above. The lead action level is not health-based. It was established based on the feasible lead level that public water systems could meet.

Maximum Contaminant Level Goal. The level of lead or copper in drinking water below which there is no known or expected risk to health. The MCLG for lead is 0 (zero) ppb, based on information from EPA and CDC that there is no identifiable level of lead that is without risk, and the MCLG for copper is 1300 ppb which is the same as the copper action level and water at or below this level is considered safe.

Explanation of the Health Effects of Lead and Copper

Lead causes serious health problems if too much enters your body from drinking water and other sources. Drinking water is just one way we consume lead. EPA estimates that less than 20% of lead exposure for the overall population is from drinking water. Infants who consume mostly mixed formula can receive 40 to 60 percent of their exposure to lead from drinking water. Other major sources include lead paint dust, soil and food, food and beverage containers, leaded gasoline and occupational exposure.

Too much lead can cause damage to the brain and kidneys, and it interferes with the production of red blood cells that carry oxygen to all parts of your body. The greatest risk of lead exposure is to infants, young children, and pregnant women. Scientists have linked the effects of lead on the brain with lowered IQ in children. Adults with kidney problems and high blood pressure can be affected by low levels of lead more than healthy adults. Lead is stored in the bones, and it can be released later in life. During pregnancy, the child receives lead from the mother's bones, which may affect brain development.

Some people who drink water containing copper in excess of the action level may, with short term exposure, experience nausea, cramps, diarrhea and/or vomiting with more severe effects associated with higher levels of copper; and with long-term exposure may experience liver or kidney damage. People with Wilson's Disease should consult their personal doctor if the amount of copper in their water exceeds action level.

How to reduce or eliminate your exposure to lead and copper in drinking water:

Lead and copper originate in plumbing materials and are released from pipes, valves and faucets by corrosion or friction. Wisconsin banned the use of lead solder in 1984. Prior to then lead was widely used in pipe solder, packing, valves, meters and as solid pipe. Brass is an alloy containing up to 15% lead, and most valves and faucets have brass bodies. Copper pipes installed prior to 1985 may have lead solder joints. Water quality affects how readily plumbing corrodes. Hard water deposits a protective scale in pipes which prevents metals from corrosion. Softened or highly treated water is more aggressive and tends to dissolve pipes and other plumbing materials. The following actions will help reduce your exposure to lead and copper in drinking water:

- Run the water until it turns cold before drinking. It is a good idea to flush the water lines before drinking any time the water has been motionless in the distribution system for 6 hours or more.
- *Use cold water for cooking and preparing baby formula.* Do not cook with or drink water from the hot water tap; lead dissolves more easily in hot water.
- *Do not boil water to remove lead.* Boiling water will not reduce lead levels.
- *Look for alternative sources or treatment of water.* If your lead result is above 15 ppb, you may want to consider purchasing bottled water or a water filter. Read the package to be sure the filter is approved to reduce lead or contact NSF International at 800-NSF-8010, or www.nsf.org for information on performance standards for water filters.
- *Identify if your plumbing fixtures contain lead.* New faucets, fittings, and valves, may contain up to 8 percent lead including those advertised or labeled as "lead-free" and may contribute lead to drinking water. Consumers should be aware of this when choosing fixtures and take appropriate precautions.

Lead and copper compliance samples represent worst-case-scenarios. You might consider taking additional water samples after flushing. Most labs charge about \$25 for a metals analysis. Take 1 sample after flushing a minute or so; and another in the middle of a day or evening, to test a random sample during periods of regular water use.

For more information call: **Steve Heraly - (920)819-2862** or visit the WDNR website to view or print a brochure about lead at <http://dnr.wi.gov/topic/drinkingwater/documents/forms/lead.pdf> and about copper at <http://dnr.wi.gov/topic/drinkingwater/documents/forms/copperdg027.pdf>